

Why Kohn-Sham and Hartree-Fock orbitals are very close to each other. Shell structure of exchange potential in atoms.

M. Cinal

Institute of Physical Chemistry of the Polish Academy of Sciences, ul. Kasprzaka 44/52, 01-224 Warszawa, Poland
(Dated: March 16, 2009)

It is found that, in closed- l -subshell atoms, the exact local exchange potential of the density functional theory is very well represented, within the region of every atomic shell, by each of the suitably shifted potentials obtained with the non-local Fock exchange operator acting on the Hartree-Fock (HF) orbitals that belong to this shell. This explains the outstanding proximity of Kohn-Sham and HF orbitals and the high quality of the Krieger-Li-Iafrate and localized HF (or, equivalently, common-energy-denominator) approximations to the exchange potential.

PACS numbers: 31.15.Ew, 71.15.Mb

Representing the quantum state of a many-electron system in terms of one-electron orbitals is simple and theoretically attractive approach. Such description is realized in the Hartree-Fock (HF) method [1], as well as in the Kohn-Sham (KS) scheme of the density-functional theory (DFT) [2] which is an efficient and robust tool, now routinely applied in the calculations of electronic properties of molecules and condensed-matters structures, even very large and complex. Though the KS scheme is formally accurate, the one-body KS potential contains the exchange-correlation (xc) potential v_{xc} , whose dependence on the electron density remains unknown. It is usually treated within the local density or generalized-gradient approximations (LDA, GGA), despite the well-known shortcomings of the LDA and GGA xc potentials (especially the self-interaction errors). Some of these deficiencies are removed when the exact form (in terms of the KS occupied orbitals) is used for the exchange part E_x of the xc energy. The exact exchange potential v_x is then found from E_x by means of the integral equation resulting from the optimized-effective-potential (OEP) approach [3(a),4,5] or by using the recently developed method based on the differential equations for the orbital shifts [6,7]. The exact potential v_x is free from self-interaction and it has correct asymptotic dependence ($-1/r$ for finite systems) at large distances r from the system; thus, unlike the HF, LDA or GGA potentials, it produces correct unoccupied states. In the DFT, the approximation, in which the exchange is included exactly but the correlation energy and potential are neglected, is known as the exchange-only KS scheme — it is applied in the present investigation. The full potential v_{xc} can also be found by means of the OEP approach when the DFT total energy includes, besides the exact E_x , the correlation energy E_c depending on all (occupied and unoccupied) KS orbitals. This makes such computation tedious, to a level undesirable in the DFT, since it involves calculating E_c with the quantum-chemistry methods, like the Møller-Plesset many-body perturbation approach.

Defined to yield the true electron density, the KS one-electron orbitals have no other direct physical meaning since they formally refer to a *fictitious* system of *non-interacting* electrons. However, it is a common practice to use these orbitals in calculations of various electronic properties; in doing

so the N -electron ground-state wave function Ψ_0 of the physical (interacting) system is approximated with the single determinant built of the KS orbitals. This approximate approach is justified by (usually) sufficient accuracy of the calculated quantities, which is close to, or often better than, that of the HF results. It seems that the success of the DFT calculations would not be possible if the KS determinant, though being formally non-physical, was not close to the HF determinant which, outside the DFT, is routinely used to approximate the wave function Ψ_0 of the *real* system. Therefore, understanding this proximity is certainly very important for the fundamentals of the DFT.

Previous calculations [8,9] have shown that, not only the whole KS and HF determinants, but also the individual occupied KS and HF orbitals $\phi_{a\sigma}(\mathbf{r})$ and $\phi_{a\sigma}^{\text{HF}}(\mathbf{r})$ in atoms are so close to each other that they are virtually indistinguishable (here the orbitals, dependent on the electron position $\mathbf{r}$ and the spin $\sigma = \downarrow, \uparrow$, are numbered with index $a = 1, \dots, N_\sigma$; $N_\downarrow + N_\uparrow = N$). This proximity is particularly remarkable for the exchange-only KS orbitals. It is surprising in view of the obvious difference between the exchange operators in the KS and HF one-electron hamiltonians (see below) and the fact that the corresponding KS and HF atomic orbital energies, $\epsilon_{a\sigma}$ and $\epsilon_{a\sigma}^{\text{HF}}$, differ substantially, up to several hartrees for core orbitals in atoms like Ar, Cu [4,5]. This apparent contradiction has not yet been resolved; in Ref. [10] it is suggested that the KS and HF determinants are close to each other “since the kinetic energy is much greater than the magnitude of the exchange energy”. The present paper investigates the proximity of the KS and HF orbitals and reveals that it results from the specific properties of the HF *orbital* exchange potentials [3]

$$v_{xa\sigma}^{\text{HF}}(\mathbf{r}) \equiv [\hat{v}_{x\sigma}^{\text{F}} \phi_{a\sigma}^{\text{HF}}(\mathbf{r})] / \phi_{a\sigma}^{\text{HF}}(\mathbf{r}). \quad (1)$$

obtained with the non-local Fock exchange operator $\hat{v}_{x\sigma}^{\text{F}}$ built of the HF orbitals. Simultaneously, it becomes clear why, in atoms, the DFT KS exact exchange potential $v_{x\sigma}(r)$ has the characteristic structure of a piecewise-like function where each part spans over the region of an atomic shell and it has distinctively different slope $dv_{x\sigma}(r)/dr$ in consecutive shells [11]. These shell constituents of $v_{x\sigma}(r)$ are found in the present Letter to be very well represented, within the re-

gion of each atomic shell, by the suitably shifted potentials $v_{\text{xa}\sigma}^{\text{HF}}(r)$, Eq. (1), for the HF orbitals $\phi_{a\sigma}^{\text{HF}}(\mathbf{r})$ that belong to this shell. Thus, it is shown how these *orbital-specific* HF potentials, which describe the exchange in the physical system, are almost directly mapped onto the KS local exchange potential $v_{\text{x}\sigma}(r)$ which is *common* to all electrons with spin σ in the KS non-interacting system. The revealed properties of $v_{\text{xa}\sigma}^{\text{HF}}(\mathbf{r})$ are also found to stand behind the high quality of the Krieger-Li-Iafrate(KLI) [3] approximation to the exact potential $v_{\text{x}\sigma}(\mathbf{r})$, as well as, of the localized HF (LHF) [12] approximation, equivalent to the common-energy-denominator approximation (CEDA) [13]. Note that the KLI potential can be derived by assuming that the exchange-only KS and HF orbitals are identical while the LHF (CEDA) potential is found when the KS and HF determinants are assumed to be equal.

The HF one-electron spin-orbitals $\phi_{a\sigma}^{\text{HF}}(\mathbf{r})$ are obtained by minimizing the mean value $\langle \Psi | \hat{H} | \Psi \rangle$ where $\hat{H}$ is the Hamiltonian of the N -electron interacting system and Ψ belongs to the subspace of normalized N -electron wave functions that are single determinants built of one-electron orbitals. Similar minimization is carried out in the (exchange-only) OEP method, but there are two additional constraints ($\sigma = \uparrow, \downarrow$) that for every trial determinant all N_σ constituent spin-orbitals $\phi_{a\sigma}(\mathbf{r})$ satisfy the KS equation with some local KS potential $v_{\text{s}\sigma}(\mathbf{r})$. The minimizing potential $v_{\text{s}\sigma}(\mathbf{r}) = v_{\text{s}\sigma}^{\text{OEP}}(\mathbf{r})$, yields, after subtracting from it the external $v_{\text{ext}}(\mathbf{r})$ and electrostatic $v_{\text{es}}(\mathbf{r})$ terms, the exact exchange potential $v_{\text{x}\sigma}(\mathbf{r}) = v_{\text{x}\sigma}^{\text{OEP}}(\mathbf{r})$. Thus, the KS equation, satisfied by the corresponding (OEP) orbitals $\phi_{a\sigma}(\mathbf{r})$ and their energies $\epsilon_{a\sigma}$, takes the form

$$\left(-\frac{1}{2}\nabla^2 + v_{\text{ext}} + v_{\text{es}}[n_{\text{tot}}] + v_{\text{x}\sigma}\right)\phi_{a\sigma} = \epsilon_{a\sigma}\phi_{a\sigma}. \quad (2)$$

The total electron density $n_{\text{tot}}(\mathbf{r}) = n_\uparrow(\mathbf{r}) + n_\downarrow(\mathbf{r})$, which enters $v_{\text{es}}[n_{\text{tot}}](\mathbf{r}) = \int d\mathbf{r}' n_{\text{tot}}(\mathbf{r}')/|\mathbf{r}' - \mathbf{r}|$, is the sum of the spin-projected densities $n_\sigma(\mathbf{r}) = \sum_{a=1}^{N_\sigma} |\phi_{a\sigma}(\mathbf{r})|^2$. In the HF equations, satisfied by the orbitals $\phi_{a\sigma}^{\text{HF}}(\mathbf{r})$ and energies $\epsilon_{a\sigma}^{\text{HF}}$, the multiplicative local exchange potential $v_{\text{x}\sigma}(\mathbf{r})$ is replaced with the non-local Fock exchange integral operator $\hat{v}_{\text{x}\sigma}^{\text{F}}(\mathbf{r})$, built of $\{\phi_{a\sigma}^{\text{HF}}\}_{a=1}^{N_\sigma}$, while the potential $v_{\text{es}}(\mathbf{r})$ is found for the HF total electron density defined in a similar way as $n_{\text{tot}}(\mathbf{r})$. Real KS and HF orbitals are used throughout this paper, dependence on σ is suppressed hereafter. Both the KS and HF equations need to be solved selfconsistently.

The exact exchange potential $v_{\text{x}} = v_{\text{x}}^{\text{OEP}}$ satisfies the OEP equation [6]

$$\sum_{a=1}^N \phi_a(\mathbf{r}) \delta\phi_a(\mathbf{r}) = 0, \quad \forall \mathbf{r}, \quad (3)$$

which results from the OEP minimization and depends on v_{x} through the orbital shifts (OS) $\delta\phi_a(\mathbf{r})$. Each OS fulfills the equation [6, 7] (below ϕ_a, ϵ_a are the solutions of Eq. (2))

$$\left(\hat{h}_s - \epsilon_a\right)\delta\phi_a(\mathbf{r}) = W_a^\perp(\mathbf{r}) \quad (4)$$

and it is subject to the constraint $\langle \phi_a | \delta\phi_a \rangle = 0$. The equation (4) includes the KS Hamiltonian $\hat{h}_s$, present in Eq. (2), and

the term (defined using the sign convention of Refs. [6, 7])

$$W_a^\perp(\mathbf{r}) = [\hat{v}_{\text{x}}^{\text{F}}(\mathbf{r}) + D_{aa} - v_{\text{x}}(\mathbf{r})] \phi_a(\mathbf{r}). \quad (5)$$

Here, we have $\hat{v}_{\text{x}}^{\text{F}} = \hat{v}_{\text{x}}^{\text{F}}[\{\phi_b\}]$, $D_{aa} = \langle \phi_a | v_{\text{x}} - \hat{v}_{\text{x}}^{\text{F}} | \phi_a \rangle$.

The OS $\delta\phi_a$ and the constant D_{aa} give, within the perturbation theory, the first-order approximations to the differences $-(\tilde{\phi}_a^{\text{HF}} - \phi_a)$ and $-(\tilde{\epsilon}_a^{\text{HF}} - \epsilon_a)$. Here, the orbitals $\tilde{\phi}_a^{\text{HF}}$ and the corresponding energies $\tilde{\epsilon}_a^{\text{HF}}$, are the solutions of the HF-like equation which is the same as Eq. (2) except for v_{x} replaced by $\hat{v}_{\text{x}}^{\text{F}}$ built of the KS orbitals $\{\phi_b\}$. The corresponding perturbation is then equal to $\delta\hat{h}_s = \hat{v}_{\text{x}}^{\text{F}} - v_{\text{x}}$ so that the first-order correction to $\phi_a(\mathbf{r})$, i.e., $-\delta\phi_a(\mathbf{r}) = -\sum_{t=1, t \neq a}^\infty c_{ta} \phi_t(\mathbf{r})$, $c_{ta} = \langle \phi_t | \delta\hat{h}_s | \phi_a \rangle / (\epsilon_t - \epsilon_a)$, satisfies Eq. (4) and the constraint indeed. Obviously, the solutions $\tilde{\phi}_a^{\text{HF}}$, $\tilde{\epsilon}_a^{\text{HF}}$ are not identical to the self-consistent HF orbitals ϕ_a^{HF} and orbital energies ϵ_a^{HF} . However, the relations $\|\Delta\phi_a - (-\delta\phi_a)\| < 0.13\|\Delta\phi_a\|$ (where $\|\phi\|^2 = \int d\mathbf{r} |\phi(\mathbf{r})|^2$), $|\Delta\epsilon_a - (-\delta\epsilon_a)| < 0.003|\Delta\epsilon_a|$ [14] found for Be and Ar, validate the representations of $\Delta\phi_a \equiv \phi_a^{\text{HF}} - \phi_a$ by $-\delta\phi_a$ and $\Delta\epsilon_a \equiv \epsilon_a^{\text{HF}} - \epsilon_a$ by $\delta\epsilon_a = -D_{aa}$ used in the presented argument; the above inequalities are obtained for $\phi_a, \epsilon_a, \delta\phi_a$ calculated as in Ref. [7], and ϕ_a^{HF} (expanded in the Slater-type-orbital basis), ϵ_a^{HF} taken from Ref. [15].

The part of $W_a(\mathbf{r}) \equiv \delta\hat{h}_s(\mathbf{r})\phi_a(\mathbf{r})$ parallel to the orbital ϕ_a is $W_a^\parallel(\mathbf{r}) = -D_{aa}\phi_a(\mathbf{r})$ and it sets the energy shift $\delta\epsilon_a$. The part $W_a^\perp(\mathbf{r}) = W_a(\mathbf{r}) - W_a^\parallel(\mathbf{r})$, perpendicular to ϕ_a , sets the OS $\delta\phi_a(\mathbf{r})$, Eq. (4). Thus, the KS and HF orbitals, $\phi_a(\mathbf{r})$, $\phi_a^{\text{HF}}(\mathbf{r})$, can be close to each other, even if the orbital energies $\epsilon_a, \epsilon_a^{\text{HF}}$, differ significantly, provided the term $W_a^\perp(\mathbf{r})$ is sufficiently small. Note that the orbitals remain unchanged when a (possibly orbital-dependent) constant is added to the Hamiltonian in the KS or HF equations.

For closed- l -subshell atom, the non-local (integral) Fock exchange operator, acting on an atomic orbital $\phi_a(\mathbf{r}) = r^{-1}\chi_{nl}(r)Y_{lm}(\theta, \varphi)$ ($a \equiv nlm$), yields $\hat{v}_{\text{x}}^{\text{F}}(\mathbf{r})\phi_a(\mathbf{r}) = r^{-1}F_{x;nl}(r)Y_{lm}(\theta, \varphi)$ where $Y_{lm}(\theta, \varphi)$ is the spherical harmonic; hereafter, n, l, m denote the principal, orbital, and magnetic quantum numbers. The factor $F_{x;nl}(r)$ is defined [1] with the exchange integrals and determines the $v_{\text{xa}}(\mathbf{r}) = v_{x;nl}(r) = F_{x;nl}(r)/\chi_{nl}(r)$. The OS $\delta\phi_a(\mathbf{r}) = r^{-1}\delta\chi_{nl}(r)Y_{lm}(\theta, \varphi)$ depends on $W_a^\perp(\mathbf{r}) = r^{-1}W_{nl}^{\perp;\text{rad}}(r)Y_{lm}(\theta, \varphi)$ through its radial part

$$W_{nl}^{\perp;\text{rad}}(r) = F_{x;nl}(r) + [D_{nl;nl} - v_{\text{x}}(r)]\chi_{nl}(r) \quad (6)$$

entering the equation for $\delta\chi_{nl}(r)$ found from Eq. (4).

Neither the OEP minimization nor the resulting OEP equation (3) imply readily that the orbital shifts $\delta\phi_a(\mathbf{r})$ are small. However, the *occupied* KS and HF orbitals differ so minutely from each other that, for atoms, the OEP total energy is only several mhartrees higher than the HF energy E_{HF} [4, 5, 16]. The proximity of individual HF and KS orbitals can be quantified with the norms $\|\delta\phi_a\|$ which are indeed very small, in comparison with $\|\phi_a\| = \|\phi_a^{\text{HF}}\| = 1$. Calculating the OS $\delta\phi_a$ with the method of Ref. [7], we obtain $\|\delta\phi_{1s}\| = 0.00669$,




FIG. 1: The OS norm square $\|\delta\phi_a\|^2$ (grey bars) and the contributions $c_{n'l;nl}^2$ (stacked bars) to it from bound states $\phi_{n'l m}$, for the occupied states $\phi_{nl m}$ in the Ar atom; the 1s bars are magnified by the factor 20.

$\|\delta\phi_{2s}\| = 0.00630$ for Be and $\|\delta\phi_a\| < 0.007$ for each occupied orbital in the Ar atom. The partition $\|\delta\phi_{nl m}\|^2 = \sum_{n' \neq n} c_{n'l;nl}^2$, plotted for Ar in Fig. 1, shows that, among the KS bound orbitals $\phi_{n'l m}$, the dominating contributions $c_{n'l;nl}^2$ to the nlm OS come from the $n'l m$ orbitals, $n' = n-1, n+1$, i.e., from the neighboring electronic shells; e.g., for $\delta\phi_{3s}$ in Ar, the largest terms $c_{n'l;nl}^2$ are found for the $n'l = 2s$ (occupied) and $n'l = 4s$ (unoccupied) orbitals. But, there remains a large part of $\|\delta\phi_{nl m}\|^2$ which cannot be attributed to higher unoccupied bound states $\phi_{n'l m}$ since the corresponding $c_{n'l;nl}^2$ terms vanish rapidly; see Fig. 1. This unaccounted part comes from *continuum* KS states ($\epsilon_{n'l} > 0$). These results confirm that the assumption $\delta\phi_a = 0$ and $\delta\phi_a - \sum_{t \neq a}^{\text{occ}} c_{ta} \phi_t = 0$, used to derive the KLI and LHF (CEDA) approximation, respectively, are very accurate but not exact.

The norms $\|\delta\phi_{nl m}\|$ have such low values because the terms $W_{nl}^{\perp; \text{rad}}(r)$ are sufficiently small for all r . This, combined with the relation

$$v_{x;nl}(r) + D_{nl;nl} = v_x(r) + \frac{W_{nl}^{\perp; \text{rad}}(r)}{\chi_{nl}(r)}, \quad (7)$$

found with Eq. (6), implies that each *shifted* KS-OEP orbital exchange potential

$$\tilde{v}_{x;nl}(r) \equiv v_{x;nl}(r) + D_{nl;nl} \quad (8)$$

is very close to the exact exchange potential $v_x(r) = v_x^{\text{OEP}}(r)$ within the r -interval $(r_{n-1,n}, r_{n,n+1}) \equiv S_n$ where the denominators in the r.h.s. of Eq. (7), i.e., the orbitals $\chi_{nl}(r)$ from the n -th atomic shell ($K, L, M, \dots$), have largest magnitudes; the shell border points $r_{n,n+1}$ for $n = 0, 1, \dots, n_{\text{occ}}$ (the respective HF points $r_{n,n+1}^{\text{HF}}$, defined precisely below, can be used) are near the positions r_n^{min} where the radial electron density $\rho(r)$ has local minima. The potentials $\tilde{v}_{x;nl}(r)$ are also close to $v_x^{\text{OEP}}(r)$ in the central parts of shells $S_{n'}$, $n' < n$, though not so tightly as for the shell S_n . The proximity of the potentials, evident in Fig. 2(a,b) for the Ar atom, holds as well for other closed- l -subshell atoms. It is disturbed in the vicinity of the nodes of $\chi_{nl}(r)$, where the potential $\tilde{v}_{x;nl}(r)$ diverges while the term $W_{nl}^{\perp; \text{rad}}(r)$ is finite and small. The potential $\tilde{v}_{x;nl}(r)$ also differs significantly from $v_x^{\text{OEP}}(r)$ within




FIG. 2: (a,b) The potentials v_x^{OEP} (solid line), $\tilde{v}_{x;nl}$ (dashed and dotted lines) in the Ar atom; the HF potentials $\tilde{v}_{x;nl}^{\text{HF}}$ follow $\tilde{v}_{x;nl}$ (within the figure resolution); (c) The differences $v_x^{\text{OEP}} - v_x^{\text{pw}}$ (dotted line), $v_x^{\text{OEP}} - v_x^{\text{KLI-HF}}$ (dashed line), $v_x^{\text{OEP}} - v_x^{\text{KLI-OEP}} = v_x^{\text{OS}}$ (solid line); the up and down arrows mark the points $r_{n,n+1}^{\text{HF}}$ and r_n^{min} , respectively.

the shells $S_{n'}$, $n' > n$, where both the functions $\chi_{nl}(r)$, $W_{nl}^{\perp; \text{rad}}(r)$ decay exponentially.

Since the KS orbitals $\phi_a^{\text{OEP}} = \phi_a[v_x^{\text{OEP}}]$ found for the exact exchange potential v_x^{OEP} are very close to ϕ_a^{HF} , the terms $F_{x;nl}(r)$, $v_{x;nl}(r)$, and $D_{nl,nl}[v_x]$ (for any v_x) obtained with $\{\phi_a^{\text{OEP}}\}$ are virtually indistinguishable from the quantities $F_{x;nl}^{\text{HF}}(r)$, $v_{x;nl}^{\text{HF}}(r)$, $D_{nl,nl}^{\text{HF}}[v_x]$ calculated with the HF orbitals $\{\phi_a^{\text{HF}}\}$. Thus, the terms $W_{nl}^{\perp; \text{rad}}[v_x, \{\phi_a^{\text{HF}}\}](r)$ are very close to $W_{nl}^{\perp; \text{rad}}[v_x, \{\phi_a^{\text{OEP}}\}](r)$. As a result, they are small for $v_x = v_x^{\text{OEP}}$, and, also, by continuity, for any potential v_x close to v_x^{OEP} . Such a class $\mathcal{V}_0$ [17] of potentials v_x that yield small $W_{nl}^{\perp; \text{rad}}[v_x, \{\phi_a^{\text{HF}}\}]$ and, consequently, lead to the KS orbitals $\phi_a[v_x]$ almost identical to ϕ_a^{HF} [18], exists provided it is possible, for given v_x , to find constants C_{nl} that make terms $U_{nl}(r) \equiv F_{x;nl}^{\text{HF}} + C_{nl}\chi_{nl}^{\text{HF}} - v_x\chi_{nl}^{\text{HF}}$ small for all r . Indeed, we then obtain $D_{nl;nl}^{\text{HF}} \approx C_{nl}$ so that the term $W_{nl}^{\perp; \text{rad}}[v_x, \{\phi_a^{\text{HF}}\}] \approx U_{nl}$ is small. Low magnitude of $U_{nl}(r)$ implies that, within each occupied shell S_n , the shifted HF potentials $\tilde{v}_{x;nl}^{\text{HF}}(r) = v_{x;nl}^{\text{HF}}(r) + C_{nl}$ ($l \in \mathcal{L}_n \equiv \{0, \dots, l_{\text{max}}^{(n)}\}$) lie very close to $v_x(r)$, and, as a result, they almost coincide with each other,

$$\tilde{v}_{x;nl}^{\text{HF}}(r) \approx \tilde{v}_{x;nl'}^{\text{HF}}(r), \quad l, l' \in \mathcal{L}_n, r \in S_n, \quad (9)$$

(similarly, as the OEP potentials $\tilde{v}_{x;nl}(r)$ do; Fig. 2(a,b)). A generalization of Eq. (9) is found when, in the expression for

$U_{nl}(r)$, the potential $v_x(r)$ is replaced by $\tilde{v}_{x;n'l'}^{\text{HF}}(r)$ for $r \in S_{n'}$ and the smallness of $U_{nl}(r)$ is accounted for. The generalized relation reads

$$F_{x;nl}^{\text{HF}} + C_{nl}\chi_{nl}^{\text{HF}} \approx (v_{x;n'l'}^{\text{HF}} + C_{n'l'})\chi_{nl}^{\text{HF}}, \quad r \in S_{n'} \quad (10)$$

and it is satisfied for suitable set of constants $\{C_{nl}\}$ for all indices (nl) , $(n'l')$ corresponding to the occupied HF orbitals. This is an *intrinsic* property of the HF orbitals (and the Fock operator), since it is *not* implied by the DFT or the definition of v_x^{OEP} , though it is revealed here by inspecting the results for $v_x = v_x^{\text{OEP}}$. The total energy $E[v_x] \equiv \langle \Psi[v_x] | \hat{H} | \Psi[v_x] \rangle$, for any $v_x \in \mathcal{V}_0$, is very close to E_{HF} (due to $\phi_a[v_x] \approx \phi_a^{\text{HF}}$), and, as a result, also to $E[v_x^{\text{OEP}}]$ since the potential v_x^{OEP} minimizes the functional $E[v_x] > E_{\text{HF}}$. This can explain why the exact exchange potential v_x^{OEP} belongs to the class $\mathcal{V}_0$ and, as a result, it gives the KS orbitals ϕ_a very close to ϕ_a^{HF} .

Assuming that the constants C_{nl} satisfying the relation (10) are known, we can construct a continuous *piecewise* potential

$$v_x^{\text{pw}}(r) = \sum_n^{\text{occ}} \theta_n^{\text{HF}}(r) v_x^{(n)}(r) \quad (11)$$

formed from the HF *shell* exchange potentials

$$v_x^{(n)}(r) \equiv \sum_{l \in \mathcal{L}_n} [v_{x;nl}^{\text{HF}}(r) + C_{nl}] \rho_{nl}^{\text{HF}}(r) / \rho_n^{\text{HF}}(r), \quad (12)$$

each applied in its shell region S_n . The points $r_{n,n+1}^{\text{HF}}$ defining the shell borders are the solutions of the continuity equation $v_x^{(n)}(r) = v_x^{(n+1)}(r)$ for $n = 1, 2, \dots, n_{\text{occ}} - 1$; $r_{0,1}^{\text{HF}} = 0$. We denote $\rho_{nl}^{\text{HF}} = 4\pi(2l+1)(\chi_{nl}^{\text{HF}})^2$, $\rho_n^{\text{HF}} = \sum_{l \in \mathcal{L}_n} \rho_{nl}^{\text{HF}}$, $\rho^{\text{HF}} = \sum_n^{\text{occ}} \rho_n^{\text{HF}}$, $\theta_n^{\text{HF}}(r) = \theta(r - r_{n-1,n}^{\text{HF}})\theta(r_{n,n+1}^{\text{HF}} - r)$. Each shell potential $v_x^{(n)}(r)$ is very close to the almost coinciding potentials $\tilde{v}_{x;nl}^{\text{HF}}(r) \equiv v_{x;nl}^{\text{HF}}(r) + C_{nl}$, $l \in \mathcal{L}_n$, for $r \in S_n$. Thus, the relation (10) with $\tilde{v}_{x;n'l'}^{\text{HF}}$ replaced by $v_x^{(n')}$ means that the terms $U_{nl}(r)$ are small for $v_x = v_x^{\text{pw}}$. This also holds for the KLI-like [3] potential

$$\check{v}_x^{\text{KLI-HF}}(r) \equiv \sum_{nl}^{\text{occ}} (F_{x;nl}^{\text{HF}} + C_{nl}\chi_{nl}^{\text{HF}}) \chi_{nl}^{\text{HF}} / \rho^{\text{HF}}, \quad (13)$$

since, due to Eq. (10), it is very close to $\tilde{v}_{x;n'l'}^{\text{HF}}(r) \approx v_x^{(n')}(r)$ for $r \in S_{n'}$, and, thus, to v_x^{pw} for all r . Therefore, both potentials v_x^{pw} and $\check{v}_x^{\text{KLI-HF}}$ belong to $\mathcal{V}_0$. In particular, it can be shown to hold for $v_x^{\text{KLI-HF}} = \check{v}_x^{\text{KLI-HF}}[\{C_{nl}^{\text{KLI-HF}}\}]$ where the constants are found from their self-consistency condition $C_{nl}^{\text{KLI-HF}} = D_{nl;nl}[v_x^{\text{KLI-HF}}, \{\chi_{n'l'}^{\text{HF}}\}]$ given in Ref. [3].

The potential $v_x^{\text{pw}}(r)$, built of the HF shell potentials $v_x^{(n)}(r)$ with $C_{nl} = C_{nl}^{\text{KLI-HF}}$, is found (see Fig. 2(c)) to be a very good approximation to the exact exchange potential $v_x^{\text{OEP}}(r)$. The quality of its approximation is almost the same as of the potentials $v_x^{\text{KLI-HF}}$ and $v_x^{\text{KLI-OEP}}$. The latter is defined similarly as $\check{v}_x^{\text{KLI-HF}}$ but with the HF orbitals $\{\chi_{nl}^{\text{HF}}\}$ replaced with the

OEP ones $\{\chi_{nl}^{\text{OEP}}\}$ and with $C_{nl} = D_{nl;nl}[v_x^{\text{OEP}}, \{\chi_{nl}^{\text{OEP}}\}]$. The potential $v_x^{\text{KLI-OEP}}$ is the dominant part of the exact exchange potential $v_x^{\text{OEP}} = v_x^{\text{KLI-OEP}} + v_x^{\text{OS}}$, where the minor part v_x^{OS} , depends linearly on $\{\delta\phi_a\}$ [6, 7]. It is this OS term $v_x^{\text{OS}}(r)$ that produces the characteristic bumps of $v_x^{\text{OEP}}(r)$ at the shell borders; cf. Fig. 2(c).

In summary, we find that when, for each HF orbital, a suitably chosen (orbital-specific) constant is added to the Fock exchange operator in the HF equation, the electrons occupying different HF orbitals are subject to very similar local exchange potentials (and hence they move in very similar total potentials) within the atomic regions where the orbital radial probability densities are substantial. This proximity is particularly tight for the exchange potentials of the orbitals that belong to the same shell and it holds in the region of this shell. Thus, each HF orbital is only very slightly disturbed when the (shifted) orbital exchange potential is replaced in the HF equation with a KS exchange potential (common to all orbitals) that, within each shell, lies very close to this orbital potential. As a result, in each shell, the KS exact exchange potential is very well approximated with the shifted HF orbital exchange potentials of this shell, and, even better, with their weighted average – the shell exchange potential, Eq. (12).

Discussions with A. Holas are gratefully acknowledged.

-
- [1] W.R. Johnson, *Atomic Structure Theory*, (Springer, Berlin, 2007).
 - [2] C. Fiolhais and F. Nogueira (ed.), *A Primer in Density Functional Theory*, Lecture Notes in Physics **620** (Springer, Berlin, 2003).
 - [3] J.B. Krieger, Y. Li, and G.J. Iafrate, Phys. Rev. A **45**, 101 (1992) (a); *ibid* **46**, 5453 (1992) (b).
 - [4] E. Engel and S.H. Vosko, Phys. Rev. A **47**, 2800 (1993).
 - [5] S. Kümmel and L. Kronik, Rev. Mod. Phys. **80**, 3 (2008).
 - [6] S. Kümmel and J.P. Perdew, Phys. Rev. Lett. **90**, 043004 (2003); Phys. Rev. B **68**, 035103 (2003).
 - [7] M. Cinal and A. Holas, Phys. Rev. A **76**, 042510 (2007).
 - [8] Q. Zhao and R.G. Parr, J. Chem. Phys. **98**, 543 (1993).
 - [9] Jiqiang Chen, R. O. Esquivel, and M. J. Stott, Philos. Mag. B, **69**, 1001(1994).
 - [10] S. Ivanov and M. Levy, J. Chem. Phys. **119**, 7087 (2003).
 - [11] O. Gritsenko, R. van Leeuwen, and E.J. Baerends, Int. J. Quant. Chem. **57**, 17 (1996).
 - [12] F. Della Sala and A. Grüning, J. Chem. Phys. **115**, 5718 (2001).
 - [13] O. V. Gritsenko and E. J. Baerends, Phys. Rev. A **64**, 042506 (2001).
 - [14] The latter is not valid for $a = \text{HOMO}$, since $D_{aa} = 0$, but $\Delta\epsilon_a = -4.13 \cdot 10^{-5}$ a.u. ($-2.645 \cdot 10^{-4}$ a.u.) for Be (Ar).
 - [15] C.F. Bunge, J.A. Barrientos and A.V. Bunge, Atomic Data and Nuclear Data Tables **53**, 113-162 (1993).
 - [16] A. Grüning and M. Ernzerhof, Phys. Rev. A **51**, 4501 (1995).
 - [17] The class $\mathcal{V}_0 = \mathcal{V}_0(\delta)$ can be defined precisely with the condition $\|\phi_a[v_x] - \phi_a^{\text{HF}}\| \leq \delta$ ($a = 1, \dots, N$) where $\delta \ll 1$.
 - [18] This results from the perturbation theory approach applied to the HF equation.